arXiv:1304.7460v2 [quant-ph] 1 May 2013

Towards loophole-free Bell inequality test with preselected unsymmetrical singlet states of light

M. Stobińska,[1, 2] F. Töppel,[3, 4] P. Sekatski,[5] and A. Buraczewski[6]

[1] *Institute of Theoretical Physics and Astrophysics,*
University of Gdańsk, ul. Wita Stwosza 57, 80-952 Gdańsk, Poland
[2] *Institute of Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland*
[3] *Max Planck Institute for the Science of Light, Guenther-Scharowsky-Str. 1/Bldg. 24, 91058 Erlangen, Germany*
[4] *Institute for Optics, Information and Photonics,*
University of Erlangen-Nürnberg, Staudtstr. 7/B2, 91058 Erlangen, Germany
[5] *Group of Applied Physics, University of Geneva,*
Chemin de Pinchat 22, CH-1211 Geneva, Switzerland
[6] *Faculty of Electronics and Information Technology,*
Warsaw University of Technology, ul. Nowowiejska 15/19, 00-665 Warsaw, Poland

The ultimate confirmation of the quantum theory by a loophole-free Bell inequality test is still missing. We examine the possibility of a loophole-free CHSH-Bell inequality test with unsymmetrical singlet states of light by employing preselection procedure which is performed prior to the test. We show an example of preselection scheme which allows to improve violation of this inequality with the micro-macro polarization singlet produced by the optimal quantum cloning. The preselection is realized by a POVM which is believed not to be useful for this purpose. The test is based on linear optical elements and is feasible within the current technology.

I. INTRODUCTION

Quantum mechanical laws apply to single particles like an electron or an atom, complex molecules involving tenths of atoms as well as to living organisms [1]. An experiment, conceptually simple and fundamental in its importance, the Bell inequalities test [2, 3], ultimately confirming or rejecting the quantum mechanical description of the world, has not yet been performed. These inequalities are violated by entangled states and reveal quantum correlations. Up to date, all experiments with light and matter suffered from either of the two loopholes: locality and detection [4]. The former is closed easily for photons with the use of fast switching polarizers. The latter however, remains an open problem, due to inefficient photon detection. It results in discarding some of the experimental data, i.e. postselection and unjustified assumption of fair data sampling.

Recently, macroscopically populated entangled states of light became available experimentally. Two kinds of them have been reported: the micro-macro polarization singlet state, where a multi-photon qubit is entangled with a single photon [5], and entangled bright squeezed vacuum, macroscopic analogs of two-photon polarization Bell states [6]. These states give hope to close the detection loophole and thus, they posed an important question of possibility of performing loophole free Bell test [7, 8]. Based on discussions in [5, 9], in Ref. [8] we emphasized the necessity of preselection for the macroscopic states in Bell tests (to improve distinguishability of the multi-photon qubit in analog detection [10]) however, without giving an example to support our claim. Moreover, later the considerations in [11] showed that all so far tested preselections are not useful for increasing the distiguishability of the micro-macro polarization singlet and analog detection.

Here, we examine a loophole-free CHSH-Bell inequality test with preselected unsymmetrical polarization singlet states of light of general form. We explicitly show an example of preselection scheme which allows to improve violation of the CHSH inequality with the micro-macro polarization singlet produced by the optimal quantum cloning.

In the unsymmetrical singlets under consideration, one of the modes is occupied by a single photon (the micro-qubit), whereas the second one contains a pair of mutually orthogonal multi-photon states (the multi-photon qubit). We assume that the average photon number in the multi-photon qubit can be controlled by some external parameter in an experimental setup, and it may vary from a single photon to the macroscopic quantity of thousands of photons. Furthermore, we consider a Bell test based only on linear optical elements. The unsymmetrical singlet is prepared before the test by a special filtering procedure applied to the mode containing the multi-photon state. The filter is described by a POVM (positive operator valued measure). For example, it may be realized by the modulus of intensity difference filter [12]. Although similar filters are believed not to be applicable for preselection [11], we find the modulus of intensity difference filter to be useful for this purpose. Filtering belongs to the conditional state preparation, not to the test. Only if the state is successfully preselected, the Bell test is performed where every measurement outcome is conclusive and is taken into account. This eliminates the necessity of data postselection and closes the detection loophole.

This paper is organized as follows. In Section II we discuss a general scenario of the CHSH-Bell inequality test with preselection strategy for an unsymmetrical singlet. Section III is devoted to a short summary of the experimentally available unsymmetrical polarization sin-



FIG. 1: Bell inequality test with unsymmetrical singlet state of light with preselection strategy.

glets of light. We further discuss the CHSH inequality violation for these states preselected by the special case of the filtering, namely the corner filter, in Section IV. Finally, we discuss the possible future steps towards the genuine loophole-free Bell test.

II. CHSH-BELL TEST WITH PRESELECTION

The Bell inequality test with an unsymmetrical singlet state and intensity measurements has to use a preselection strategy [8, 10]. We call the singlet unsymmetrical if the dimensions of the Hilbert spaces corresponding to its modes are unequal. Let us start our discussion with the two-mode unsymmetrical polarization singlet state of a general form

$$|\Psi^-\rangle = 1/\sqrt{2}(|1_\varphi\rangle_A|\bar{\Phi}\rangle_B - |1_{\varphi_\perp}\rangle_A|\Phi\rangle_B) \tag{1}$$

and an arbitrary preselection strategy executed by a POVM $\mathcal{P}$. The states $|1_\varphi\rangle$ and $|1_{\varphi_\perp}\rangle$ denote the micro-qubit, e.g. a single photon in polarization φ and $\varphi_\perp$ respectively, whereas $|\Phi\rangle$ and $|\bar{\Phi}\rangle$ are multi-photon states which constitute the multi-photon qubit. Of course, the two states of the qubits are pairwise orthogonal $\langle 1_\varphi|1_{\varphi_\perp}\rangle = 0$, $\langle\Phi|\bar{\Phi}\rangle = 0$.

The setup for the Bell test is depicted in Fig. 1. The multi-photon part of the state (mode B) impinges on a beam splitter (BS) with a low reflectivity r, e.g. 10%, which taps only a small fraction of the state leaving it almost unaffected. Next, the preselection is performed on the reflected part: it is filtered by the POVM $\mathcal{P}$ and the result is feed forwarded to the transmitted beam. This procedure belongs to the conditional state preparation before the test. After the successful preselection, the Bell test consists of polarization rotations, by angles α, α' on mode A and β, β' on mode B, and intensity measurements of polarization components of both modes.

In general, the operator $\mathcal{P}$ may suffer from lack of the rotational invariance being an important property of the original singlet and thus, the form of the preselected state may be basis dependent. For example, for the modulus of intensity difference filter it is known that it improves the distinguishability of a multi-photon state for a measurement in one polarization basis but deteriorates for measurements in other polarization bases. Thus, such filtering strategies are believed to be useless for preselection [11]. This problem arises if the usual settings for the CHSH-Bell inequality are considered: $\alpha = \pi/4$, $\alpha' = -\pi/4$, $\beta = 0$, $\beta' = \pi/2$. However, it does not need to be the case if nonorthogonal polarization directions in measurements are chosen, at the expense of lower Bell inequality violation. Moreover, the distinguishability is not the only factor contributing to the Bell parameter. We will show that in fact, for a wise choice of the rotation angles distinguishability measured in the basis unbiased with respect to the basis in which we write the original superposition does not contribute.

We consider the CHSH inequality with Bell parameter

$$B = E(\alpha,\beta) + E(\alpha,\beta') + E(\alpha',\beta) - E(\alpha',\beta') \tag{2}$$

and $E(\alpha,\beta) = \langle\mathcal{O}^{(m)}(\alpha)\otimes\mathcal{O}^{(M)}(\beta)\rangle$ is the correlation function where one observer, Alice, measures the microscopic part (mode A) and the other, Bob, measures the multi-photon part of the singlet (mode B). We assume the ideal measurement operator $\mathcal{O}^{(m)}(\alpha) = |1_\alpha\rangle\langle 1_\alpha| - |1_{\alpha_\perp}\rangle\langle 1_{\alpha_\perp}|$ for the microscopic qubit, while for the multi-photon one we take the binary threshold detection operator $\mathcal{O}^{(M)}(\beta)$ adapted to the preselection strategy $\mathcal{P}$. The value $+1$ (-1) is assign to this observable when the state $|\Phi\rangle$ $(|\bar{\Phi}\rangle)$ is identified. Rotating the polarization of the microscopic part by an angle α yields

$$\begin{aligned} |1_\alpha\rangle &= \cos(\alpha/2)\,|1\rangle + \sin(\alpha/2)\,|1_\perp\rangle\,, \\ |1_{\alpha_\perp}\rangle &= -\sin(\alpha/2)\,|1\rangle + \cos(\alpha/2)\,|1_\perp\rangle\,, \end{aligned}$$

which allows expressing the micro observable in terms of the projectors in the reference basis $\varphi = 0$

$$\begin{aligned} \mathcal{O}^{(m)}(\alpha) = &\cos\alpha\,(|1\rangle\langle 1| - |1_\perp\rangle\langle 1_\perp|) \\ &+ \sin\alpha\,(|1\rangle\langle 1_\perp| + |1_\perp\rangle\langle 1|)\,. \end{aligned} \tag{3}$$

After short algebra, we obtain the correlation function for the state in Eq. (1)

$$E(\alpha,\beta) = -\cos\alpha\,V^\theta(\beta) - \sin\alpha\,A^\theta(\beta), \tag{4}$$

where $V^\theta(\beta) = \langle\Phi^\theta|\mathcal{O}^{(M)}(\beta)|\Phi^\theta\rangle$ is called the visibility and $A^\theta(\beta) = \langle\bar{\Phi}^\theta|\mathcal{O}^{(M)}(\beta)|\Phi^\theta\rangle$ is denoted as the antivisibility of the state Φ preselected in the polarization basis θ and observed in the polarization basis β. In this derivation we took that $V^\theta_\perp(\beta) = \langle\bar{\Phi}^\theta|\mathcal{O}^{(M)}(\beta)|\bar{\Phi}^\theta\rangle = -V^\theta(\beta)$ and that $A^\theta(\beta)$ is real-valued. The antivisibility quantifies the ability of the observable $\mathcal{O}^{(M)}(\beta)$ to erase the information on which state of $|\bar{\Phi}_\theta\rangle$ or $|\Phi_\theta\rangle$ entered the detector. We obtain a general formula for the Bell parameter after inserting Eq. (4) into Eq. (2)

$$\begin{aligned} B = &-\cos\alpha\,(V^\theta(\beta) + V^\theta(\beta')) - \sin\alpha\,(A^\theta(\beta) + A^\theta(\beta')) \\ &- \cos\alpha'\,(V^\theta(\beta) - V^\theta(\beta')) - \sin\alpha'\,(A^\theta(\beta) - A^\theta(\beta')). \end{aligned}$$

We further optimize the Bell parameter with respect to the rotations on mode A and B by choosing $\theta = 0$, $\alpha' = -\alpha$, $\beta' = \beta + \pi/2$, see Fig. 2. We get

$$B_{opt} = -2\cos\alpha\,V(\beta) - 2\sin\alpha\,A(\beta + \pi/2).$$



FIG. 2: The usual (red and blue) and optimal (red and black) angle settings for CHSH inequality test.

Next, to get its maximal value we look for the α which fulfills the condition of the extremum $\partial_\alpha B_{opt} = 0$ and find

$$\begin{aligned} \cos\alpha &= \sqrt{V^2(\beta)/(V^2(\beta) + A^2(\beta+\pi/2))}, \\ \sin\alpha &= \sqrt{A^2(\beta+\pi/2)/(V^2(\beta) + A^2(\beta+\pi/2))}. \end{aligned} \tag{5}$$

Additionally, we chose $\beta = 0$ and obtain

$$B_{opt} = 2\sqrt{V^2(0) + A^2(\pi/2)}. \tag{6}$$

The final form of the Bell parameter depends on two factors: the visibility $V(0) = V$ measured in the reference basis and the antivisibility $A(\pi/2) = A$ measured in the basis unbiased with respect to the reference one. This holds true for any preselection strategy and any unsymmetrical singlet state of the form of Eq. (1). A violation of Bell's inequality is obtained if $V^2(0) + A^2(\pi/2) > 1$. The non-orthogonal choice for the measurement settings on the micro side, Eq. (5), indicates that the unsymmetrical singlet after preselection loses the phase-covariant symmetry, and the best correlations are observed when measuring both modes in the $\beta = 0$ basis. If instead of taking the preselected unsymmetrical multi-photon singlet we will take the usual two-photon singlet, so that there will be a single photon in mode B, we will obtain $V(0) = A(\pi/2) = 1$ which gives $\alpha = \pi/4$. Thus, we will restore the usual CHSH orthogonal settings. The result in Eq. (6) can be easily generalized for $V^\theta_\perp(\beta) \neq -V^\theta(\beta)$ and complex $A^\theta(\beta)$: $B_{opt} = \sqrt{(V_\perp(0) - V(0))^2 + (A(\pi/2) + A^*(\pi/2))^2}$.

III. EXAMPLE: UNSYMMETRICAL POLARIZATION SINGLET STATES OF LIGHT EMERGING FROM PHASE-COVARIANT QUANTUM CLONER

In this section we will discuss a specific example of the experimentally available unsymmetrical polarization singlet states of light. They are produced in the process of the phase-covariant optimal quantum cloning. It is based on phase sensitive parametric amplification [13–15] and requires a pair of linearly polarized photons in a standard singlet state, obtained through parametric down conversion, as input. The seeding single photon is coherently amplified to produce a multi-photon state, by an intensely pumped high gain g nonlinear medium (the cloner). The equatorial states of the Poincaré sphere of all polarization states, parametrized by the polar angle $\varphi \in \langle 0, 2\pi)$, are privileged for the phase-covariant cloners, since only for this subspace their Hamiltonian $H = \frac{i\chi}{2}\left({a^\dagger_\varphi}^2 + {a^\dagger_{\varphi_\perp}}^2\right) + \text{h.c.}$ is rotationally invariant and they work equally well for all of them. The operators $a^\dagger_\varphi$ and $a^\dagger_{\varphi_\perp}$ denote the creation operators for the equatorial polarization modes φ and $\varphi_\perp$, respectively and χ is the coupling strength, proportional to the pumping power. We restrict ourselves to the equatorial polarization state subspace for the seeding photon. The subspace basis is set by two states, $|1_\varphi\rangle = 1/\sqrt{2}(|1_H\rangle + e^{i\varphi}|1_V\rangle)$ and its orthogonal counterpart $|1_{\varphi_\perp}\rangle$, where $|k_H\rangle$ ($|l_V\rangle$) denote k (l) photons polarized horizontally (vertically) and $\varphi_\perp = \varphi + \pi$. Due to its rotational invariance, we express the initial singlet in this basis $|\psi^-\rangle = 1/\sqrt{2}(|1_\varphi\rangle_A|1_{\varphi_\perp}\rangle_B - |1_{\varphi_\perp}\rangle_A|1_\varphi\rangle_B)$. Cloning is a unitary process and the original two-photon entanglement is transferred to a unsymmetrical singlet with

$$\begin{aligned} |\Phi\rangle &= \sum_{i,j=0}^{\infty} \gamma_{ij} \big|(2i+1)_\varphi, (2j)_{\varphi_\perp}\big\rangle, \\ |\bar\Phi\rangle &= \sum_{i,j=0}^{\infty} \gamma_{ij} \big|(2j)_\varphi, (2i+1)_{\varphi_\perp}\big\rangle. \end{aligned} \tag{7}$$

$|\Phi\rangle$ and $|\bar\Phi\rangle$ are the amplified single photons, with the real-valued probability amplitude $\gamma_{ij} = \cosh^{-2} g\,(\tanh g/2)^{i+j}\,\sqrt{(1+2i)!(2j)!}/i!/j!$. In the experiment, their average population varied from less than one up to 10^4 of photons, depending on the parametric gain g. Due to different parity of occupation number in the Fock state basis, $|\Phi\rangle$ and $|\bar\Phi\rangle$ are orthogonal. However, in high photon number regime photon detectors are not single photon resolving [15] and distinguishability of the multi-photon states is quite low, making them inapplicable for quantum protocols and Bell inequality test.

Distinguishability of the multi-photon qubit can be improved by quantum state filtering performed by certain POVM filters. They modify the state but preserve quantum superpositions. Recently, such a filter has been proposed [12]: the modulus of intensity difference filter selects two-mode states of light whose mode populations differ by more than a certain threshold δ_{th}. It estimates the absolute value of difference instead of the difference. It does not provide any information on which polarization mode was more populated and thus, it is not able to distinguish the multi-photon states and preserves superpositions. It performs almost a nondestructive measurement implemented by the tapping and feed forward loop, i.e. the filtered output state is almost pure. Qualitatively, it approximates the following operation

$$\mathcal{P}_{\mathrm{MDF}} = \sum_{\substack{k,l=0 \\ |k-l| \geq \delta_{th}}}^{\infty} |k,l\rangle\langle k,l|, \tag{8}$$

where $|k,l\rangle$ is a polarization two mode Fock state. Its physical implementation and action on a multi-photon states is examined in detail in [12]. In fact, the suggested physical implementation of the modulus of intensity difference filter allows to impose an arbitrary filtering condition on combination of the sum $\sigma = k + l$ and the difference $\Delta = k - l$ of photon occupations in the two polarization modes.

IV. CHSH-BELL TEST FOR THE UNSYMMETRICAL POLARIZATION SINGLET EMERGING FROM PHASE-COVARIANT QUANTUM CLONER

In order to investigate the Bell inequality violation for the unsymmetrical singlet described in Section III, we re-write the multi-photon states given in Eq. (7) as superpositions of states of a fixed photon number $2k+1$ distributed over the two polarization modes

$$|\Phi\rangle = \sum_{k=0}^{\infty} \beta_k \, |\Phi_k\rangle, |\Phi_k\rangle = \tfrac{1}{\sqrt{\mathcal{N}_k}} \left(a^{\dagger\,2} + a_{\perp}^{\dagger\;2}\right)^k a^{\dagger} \, |0\rangle, \tag{9}$$

$$|\bar{\Phi}\rangle = \sum_{k=0}^{\infty} \beta_k |\Phi_{\perp\,k}\rangle, |\Phi_{\perp\,k}\rangle = \tfrac{1}{\sqrt{\mathcal{N}_k}} \, \left(a^{\dagger\,2} + a_{\perp}^{\dagger\;2}\right)^k a_{\perp}^{\dagger} \, |0\rangle,$$

where we took $\varphi = 0$, $\mathcal{N}_k = 4^k \, k!^2 \, (1+k)$,

$$\beta_k = \cosh^{-2} g \, (\tanh g)^k \, \sqrt{1+k} \tag{10}$$

and $\sum_{k=0}^{\infty} \beta_k^2 = 1$.

A. No preselection

Since the states are given by a convex sum of different photon number sectors, we note that the visibility, the antivisibility and the Bell parameter can also be expressed in this form

$$V = \sum_{k=0}^{\infty} \beta_k^2 \, V_k, \quad A = \sum_{k=0}^{\infty} \beta_k^2 \, A_k, \quad B = \sum_{k=0}^{\infty} \beta_k^2 \, B_k, \tag{11}$$

with $B_k = 2\sqrt{(V_k)^2 + (A_k)^2}$. Here, the sum components V_k, A_k and B_k are the visibility, antivisibility and the Bell parameter computed for the photon number sector $|\Phi_k\rangle$ and $|\Phi_{\perp\,k}\rangle$. This observation significantly simplifies the analysis of the multi-photon states. In the following subsection, we will show that the decomposition in Eq. (11) holds true also in case of preselected multi-photon qubits. In order to analyze the difficulties with Bell inequality violation we adapt the following observable

$$\mathcal{O}^{(M)}(\beta) = \left(\sum_{\substack{k,l=0 \\ k-l \geq 0}}^{\infty} - \sum_{\substack{k,l=0 \\ k-l<0}}^{\infty} \right) |k_\beta, l_{\beta_\perp}\rangle\langle k_\beta, l_{\beta_\perp}|. \tag{12}$$

It is well-suited to the photon number distribution of the multi-photon qubit in Eq. (7): $|\Phi\rangle$ and $|\bar{\Phi}\rangle$ have unequal average population in the two polarization modes. This property allows to distinguish these states in analog detection. The visibility and antivisibility evaluated for this observable equal

$$V_k = \tfrac{1}{\mathcal{N}_k} \sum_{j=0}^{k} \binom{k}{j}^2 (2j+1)! \, (2k-2j)! \, \mathrm{sgn}(4j+1-2k), \tag{13}$$

$$\begin{aligned} A_k = & \tfrac{1}{\mathcal{N}_k} \tfrac{1}{2^{2k+1}} \sum_{j=0}^{k} \sum_{p=0}^{2j+1} \sum_{q=0}^{2k-2j} \sum_{j'=0}^{k} \sum_{p'=0}^{2j'} \sum_{q'=0}^{2k+1-2j'} \\ & \binom{k}{j} \binom{2j+1}{p} \binom{2k-2j}{q} (-1)^q \\ & (p+q)! \, (2k+1-p-q)! \, \mathrm{sgn}(2(p+q)-2k-1) \\ & \binom{k}{j'} \binom{2j'}{p'} \binom{2k+1-2j'}{q'} (-1)^{1-q'} \, \delta_{p+q=p'+q'} . \end{aligned} \tag{14}$$

In Fig. 3 we depicted the visibility, the antivisibility and the Bell parameter for each $2k+1$-photon-number sector separately. We note that V_k and A_k both quickly tend to the value less than $1/\sqrt{2}$ for all k and the Bell parameter B_k drops below 2 for $k \geq 3$. The probability distribution β_k^2 depends on the amplification gain g and it gives a weight to each B_k contributing to the Bell parameter B. Since the greatest weight is given to the small values of k, see Fig. 4, it is possible to weakly violate the CHSH-Bell inequality for a very small average photon population, so for very small g. Note that the considered state approaches the Bell singlet state in the limit $g \to 0$. For $g = 0.8$ $B = 2.06$, but for $g = 1.1$ we found $B = 2.01$. Thus, Bell inequality violation for the multi-photon unsymmetrical singlet with the multi-photon qubit given in Eq. (7) and the population of already few photons on average, is impossible.

B. General preselection strategy

A preselection strategy described by a POVM of the form

$$\mathcal{P}_{\mathrm{C}} = \sum_{\substack{n,m=0 \\ \mathrm{C}(\sigma=n+m, \Delta=n-m)}}^{\infty} |n,m\rangle\langle n,m|, \tag{15}$$

with a general preselection condition on σ and Δ, insignificantly influences the form of the convex sum in



FIG. 3: Plot of the visibility V_k, antivisibility A_k (top figure) and the Bell parameter B_k (bottom figure) given in Eq. (11) computed as a function of k corresponding to $(2k+1)$th photon number sector of the multi-photon states with no preselection ($|\Phi_k\rangle$ and $|\Phi_{\perp\,k}\rangle$).



FIG. 4: Plot of β_k^2 defined in Eq. (10) for $g = 0.8$ (dot) and $g = 1.1$ (cross).

which the Bell parameter is expressed in Eq (11). The detailed calculations are presented in Appendix A. Assuming that the condition $\mathrm{C}(\sigma, \Delta)$ is symmetric, i.e. $\mathrm{C}(\sigma, \Delta) = \mathrm{C}(\sigma, -\Delta)$, we show that the preselection modifies both multi-photon states in the same way

$$|\Phi_k^P\rangle = \frac{1}{\sqrt{\mathcal{N}_k^P}}\,\mathcal{P}_{\mathrm{C}}\,|\Phi_k\rangle,\ |\bar{\Phi}_k^P\rangle = \frac{1}{\sqrt{\mathcal{N}_k^P}}\,\mathcal{P}_{\mathrm{C}}\,|\bar{\Phi}_k\rangle, \qquad (16)$$

$$|\Phi^P\rangle = \sum_{k=0}^{\infty}\beta_k^P\,|\Phi_k^P\rangle,\ |\bar{\Phi}^P\rangle = \sum_{k=0}^{\infty}\beta_k^P|\bar{\Phi}_k^P\rangle,$$

where $\mathcal{N}_k^P$ (Eq. 20) is the new normalization constant and β_k^P (Eq. (23)) is the new probability amplitude in the decomposition of the states into the fixed-number-of-photons sectors. The straightforward calculation shows that the visibility equals $V^P = \langle\Phi^P|\mathcal{O}^{(M)}(0)|\Phi^P\rangle = \sum_{k=0}^{\infty}\left(\beta_k^P\right)^2 V_k^P$ and antivisibility $A^P = \langle\Phi^P|\mathcal{O}^{(M)}(\frac{\pi}{2})|\bar{\Phi}^P\rangle = \sum_k^{\infty}\left(\beta_k^P\right)^2 A_k^P$ with V_k^P given by Eq. (13) and A_k^P given by Eq. (14) where $\sum_{j=0}^{k}$ is replaced with $\sum_{j=0;\mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}^{k}$, $\sum_{j'=0}^{k}$ with $\sum_{j=0;\mathrm{C}(\sigma=2k+1,\Delta=4j'-1-2k)}^{k}$ and the normalization constant $\mathcal{N}_k$ with $\mathcal{N}_k^P$. For these quantities the Bell parameter keeps its previous form $B = \sum_{k=0}^{\infty}\left(\beta_k^P\right)^2 B_k^P$ with $B_k^P = 2\sqrt{\left(V_k^P\right)^2 + \left(A_k^P\right)^2}$.

C. Example: the corner filter

We will examine now the preselection procedure $\mathcal{P}_+$ for the following preselection condition $\mathrm{C}(\sigma, \Delta)$: $\sigma - \delta_{th} \leq |\Delta|$, where δ_{th} is a threshold. This condition means that those components of the multi-photon superpositions are selected, whose polarization modes population difference is higher than the population sum reduced by δ_{th}. It is clear that the filtering is most restrictive if $\delta_{th} = 0$. Here $|\Delta| = \sigma$ and the N00N-like components are filtered out from the polarization singlet. The case of $\delta_{th} \to \infty$ corresponds to no filtering since $0 \leq |\Delta|$ is always fulfilled. We call this kind of filtering the corner filter, since the analysis of the modification of a state in terms of its photon number distribution shows that the filter preserves these components of a superposition which belong to the region in the shape of a corner. Photon number distributions for the original multi-photon states and for the ones preselected with the corner filter are depicted in Fig. 5.

Fig. 6 depicts the visibility V_k^P, antivisibility A_k^P and Bell parameter B_k^P evaluated for the preselected with the corner filter multi-photon states with threshold values $\delta_{th} = 0$ (dots) and $\delta_{th} = 2$ (crosses). For $\delta_{th} = 0$ the states $|\Phi^P\rangle$ and $|\bar{\Phi}^P\rangle$ become superpositions of $|2i{+}1, 0_\perp\rangle$ and $|0, (2i+1)_\perp\rangle$ states respectively, with varying i. The preselected polarization singlet from Eq. (1) takes the new form of a superposition of polarization $N11N$ states with odd N

$$|\Psi_P^-\rangle = \sum_{i=0}^{\infty}\tilde{\gamma}_{i0}\big(|1\rangle_A|(2i+1)_\perp\rangle_B - |1_\perp\rangle_A|2i+1\rangle_B\big), \quad (17)$$

where $N = 2i + 1$ and $\tilde{\gamma}_{i0} = \sqrt{\cosh g/2}\gamma_{i0}$. In this case, V_k^P gets its maximal value 1 for all k. However, A_k^P decreases to 0 for large k. Thus, we still see decreasing tendency for the B_k^P with increasing k, but after preselection $B_k^P > 2$ for all k. Thus, for all g the Bell parameter



FIG. 5: Photon number distribution of multi-photon states (a) $|\Phi\rangle$, (b) $|\bar{\Phi}\rangle$ before and after preselection (d) $|\Phi^P\rangle$ & $|\bar{\Phi}^P\rangle$ with the corner filter. Action of the filter in the photon number space is shown in (c): only the components of filtered superposition which belong to the blue area are preserved.

exceeds the classical value. For $g = 0.8$ $B = 2.44$. Since β_k^P keeps its maximum at $k = 0$, the violation decreases and asymptotically $B \to 2$ for increasing g.

For $\delta_{th} = 2$ in turn, the states $|\Phi^P\rangle$ and $|\bar{\Phi}^P\rangle$ are superpositions not only of $|2i+1, 0_\perp\rangle$ and $|0, (2i+1)_\perp\rangle$ so $N00N$s with odd N, but also include terms like $|1, (2j)_\perp\rangle$ and $|2j, 1_\perp\rangle$, i.e. superpositions of $N11N$ states with even N. The antivisibility decreases slower with k however, but the visibility achieves its maximum value only for large k. Thus, we obtain smaller value for the Bell parameter. For $g = 0.8$ $B = 2.13$.

V. CONCLUSIONS

We have discussed a possibility of performing a loophole free Bell test with the use of multi-photon quantum states of light and preselection. For this purpose, we examined the CHSH inequality and exemplary unsymmetrical polarization singlet state with the multi-photon states produced in optimal phase covariant quantum cloning. Our work is a proof-of-principle: we show that it is possible to apply a feasible quantum state engineering to multi-photon states and in this way to overcome the problem of inefficient analog detection and violate the classical bound. The quantum engineering is performed by a POVM, which prepares the states to the test. It belongs to the conditional state preparation, not to the test. The exact form of the POVM should be adapted to the states entering the test. For the states we discussed, the corner filter is a good choice. It filters out the N00N-like components from the initial superpositions.

We do not claim that the discussed multi-photon states and the CHSH Bell inequality are the best strategy for



FIG. 6: Plot of the visibility V_k^P (top figure), antivisibility A_k^P (middle figure), and the Bell parameter B_k^P (bottom figure) computed as a function of k corresponding to $(2k+1)$th photon number sector of the preselected multi-photon states ($|\Phi_k^P\rangle$ and $|\Phi_{\perp\, k}^P\rangle$) with the corner filter for $\delta = 0$ (dot) and $\delta = 2$ (cross). For $g = 0.8$ and $\delta = 0$ total $B = 2.44$, whereas for $\delta = 2$ $B = 2.13$.

obtaining the loophole free violation. We believe however, that the technique of preselection we present here is worth exploring and may be useful in the context of Bell inequalities with analog detection.

It is also worth noting that so far there is no proposal allowing for a genuine macroscopic violation of a Bell inequality. From Figs. 3 and 6 it is clear that the photon number sectors which contribute to the violation most, come from the small photon numbers. The Bell parameter decreases with increasing photon number $2k+1$. This is a general tendency one observes also for the bright squeezed vacuum state and other Bell inequalities, with observables which are dichotomic or not. Indeed, the preselection helps to increase the values of B for all k but the question what is the observable which will reverse the decreasing trend presented in these figures, so that the high photon numbers were contributing to violation most, remains open.

Acknowledgments

This work is supported by the EU 7FP Marie Curie Career Integration Grant No. 322150 "QCAT", NCN grant No. 2012/04/M/ST2/00789, FNP Homing Plus project No. HOMING PLUS/2012-5/12 and MNiSW co-financed international project No. 2586/7.PR/2012/2. Computations were carried out at the CI TASK in Gdańsk and Cyfronet in Kraków.

Appendix A

The preselection modifies the sectors of the fixed photon number states in the following way

$$\begin{aligned} |\Phi_k^P\rangle =& \frac{1}{\sqrt{\mathcal{N}_k^P}}\,\mathcal{P}_{\mathrm{C}}\,|\Phi_k\rangle \qquad (18)\\ =& \frac{1}{\sqrt{\mathcal{N}_k^P}} \sum_{\substack{j=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{k} \binom{k}{j} \sqrt{(2j+1)!\,(2k-2j)!}\\ & |2j+1,2k-2j\rangle,\\ |\bar{\Phi}_k^P\rangle =& \mathcal{P}_{\mathrm{C}}\,|\bar{\Phi}_k\rangle \qquad (19)\\ =& \frac{1}{\sqrt{\bar{\mathcal{N}}_k^P}} \sum_{\substack{j=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4j-1-2k)}}^{k} \binom{k}{j} \sqrt{(2j)!\,(2k+1-2j)!}\\ & |2j,2k+1-2j\rangle, \end{aligned}$$

with normalization constants equal

$$\mathcal{N}_k^P = \sum_{\substack{j=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{k} \binom{k}{j}^2 (2j+1)!\,(2k-2j)!, \qquad (20)$$

$$\bar{\mathcal{N}}_k^P = \sum_{\substack{j=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4j-1-2k)}}^{k} \binom{k}{j}^2 (2j)!\,(2k+1-2j)!. \qquad (21)$$

The multi-photon states equal

$$\begin{aligned} |\Phi^P\rangle =& \sum_{k=0}^{\infty} \beta_k^P\,|\Phi_k^P\rangle, \qquad (22)\\ \beta_k^P =& C_g^{-2}\left(\frac{T_g}{2}\right)^k \frac{1}{k!}\sqrt{\frac{\mathcal{N}_k^P}{\mathcal{N}}}, \quad \sum_{k=0}^{\infty}\left(\beta_k^P\right)^2 = 1, \qquad (23)\\ \mathcal{N} =& C_g^{-4}\sum_{k=0}^{\infty}\left(\frac{T_g}{2}\right)^{2k} \frac{1}{k!^2}\,\mathcal{N}_k^P, \qquad (24) \end{aligned}$$

$$\begin{aligned} |\bar{\Phi}^P\rangle =& \sum_{k=0}^{\infty} \bar{\beta}_k^P|\bar{\Phi}_k^P\rangle, \qquad (25)\\ \bar{\beta}_k^P =& C_g^{-2}\left(\frac{T_g}{2}\right)^k \frac{1}{k!}\sqrt{\frac{\bar{\mathcal{N}}_k^P}{\bar{\mathcal{N}}}}, \quad \sum_{k=0}^{\infty}\left(\bar{\beta}_k^P\right)^2 = 1, \qquad (26)\\ \bar{\mathcal{N}} =& C_g^{-4}\sum_{k=0}^{\infty}\left(\frac{T_g}{2}\right)^{2k} \frac{1}{k!^2}\,\bar{\mathcal{N}}_k^P, \qquad (27) \end{aligned}$$

Let's change the variable j to j' in $\bar{\mathcal{N}}_k^P$, so $j' = k-j$ ($j = k-j'$). The sum over j' remains from 0 to k.

$$\begin{aligned} \bar{\mathcal{N}}_k^P =& \sum_{\substack{j'=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4(k-j')-2k-1)}}^{k} \binom{k}{k-j'}^2 (2(k-j'))!\,(2k+1-2(k-j'))!\\ =& \sum_{\substack{j'=0\\ \mathrm{C}(\sigma=2k+1,\Delta=-(4j'+1-2k))}}^{k} \binom{k}{j'}^2 (2k-2j')!\,(2j'+1)!. \qquad (28) \end{aligned}$$

Assuming that $\mathrm{C}(\sigma,\Delta)$ is symmetric with respect to Δ, we got $\mathrm{C}(\sigma=2k+1,\Delta=-(4j+1-2k)) = \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)$, so $\bar{\mathcal{N}}_k^P = \mathcal{N}_k^P$ and therefore $\bar{\mathcal{N}} = \mathcal{N}$ and $\bar{\beta}_k^P = \beta_k^P$.

Visibility takes the form

$$\begin{aligned} V^P =& \sum_{k,k'=0}^{\infty} \beta_k^P\,\beta_{k'}^P\,\langle\Phi_k^P|\mathcal{O}^{(M)}(0)|\Phi_{k'}^P\rangle \qquad (29)\\ =& \sum_{k}^{\infty}\left(\beta_k^P\right)^2\,\langle\Phi_k^P|\mathcal{O}^{(M)}(0)|\Phi_k^P\rangle\\ =& \sum_{k=0}^{\infty}\left(\beta_k^P\right)^2\,V_k^P \end{aligned}$$

with

$$V_k^P = \frac{1}{\mathcal{N}_k^P} \sum_{\substack{j=0\\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{k} \binom{k}{j}^2 (2j+1)!\,(2k-2j)!\,\mathrm{sgn}(4j+1-2k). \qquad (30)$$

We note that the form of V^P simplifies due to the orthogonality condition

$$\begin{gathered} \langle\Phi_k^P|\mathcal{O}^{(M)}(0)|\Phi_{k'}^P\rangle \sim\\ \langle 2j'+1,2k'-2j'|\tilde{n},\tilde{m}\rangle\langle\tilde{n},\tilde{m}|2j+1,2k-2j\rangle\\ \downarrow\\ j=j', k=k'. \end{gathered}$$

Similar considerations (see below) allow to show that the antivisibility equals

$$\begin{aligned} A^P &= \sum_{k,k'=0}^{\infty} \beta_k^P \, \bar{\beta}_{k'}^P \, \langle \Phi_k^P | \mathcal{O}^{(M)}(\tfrac{\pi}{2}) | \bar{\Phi}_{k'}^P \rangle \qquad (31) \\ &= \sum_k^{\infty} \beta_k^P \, \bar{\beta}_k^P \, \langle \Phi_k^P | \mathcal{O}^{(M)}(\tfrac{\pi}{2}) | \bar{\Phi}_k^P \rangle \\ &= \sum_k^{\infty} \left(\beta_k^P\right)^2 \, A_k^P . \end{aligned}$$

In order to evaluate the mean value of the observable $\mathcal{O}^{(M)}(\frac{\pi}{2})$ expressed in the basis complementary to the reference basis where the multi-photon states are given

$$|\Phi_k^P\rangle = \tfrac{1}{\sqrt{\mathcal{N}_k^P}} \sum_{\substack{j=0 \\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{\infty} \binom{k}{j} \, a^{\dagger\, 2j+1} \; \bar{a}^{\dagger\, 2k-2j} \; |0\rangle,$$

we perform transformation of the annihilation operators which corresponds to polarization rotation

$$a^\dagger = \frac{a_\gamma^\dagger + \bar{a}_\gamma^\dagger}{\sqrt{2}}, \qquad \bar{a}^\dagger = \frac{a_\gamma^\dagger - \bar{a}_\gamma^\dagger}{\sqrt{2}}, \qquad \gamma \equiv -\tfrac{\pi}{2}. \qquad (32)$$

The operators $a^\dagger$ and $\bar{a}^\dagger$ correspond to the reference basis whereas $a_\gamma^\dagger$ and $\bar{a}_\gamma^\dagger$ correspond the the basis rotated by angle γ. The rotation amounts to

$$\begin{aligned} a^{\dagger\, 2j+1} \; \bar{a}^{\dagger\, 2k-2j} &= \frac{\left(a_\gamma^\dagger + \bar{a}_\gamma^\dagger\right)^{2j+1}}{\sqrt{2}^{2j+1}} \frac{\left(a_\gamma^\dagger - \bar{a}_\gamma^\dagger\right)^{2k-2j}}{\sqrt{2}^{2k-2j}} \\ &= \tfrac{1}{\sqrt{2}^{2k+1}} \sum_{p=0}^{2j+1} \sum_{q=0}^{2k-2j} \binom{2j+1}{p} \binom{2k-2j}{q} \\ &\quad a^{\dagger\, p} \; \bar{a}^{\dagger\, 2j+1-p} \; (-1)^{2k-2j-q} \; a^{\dagger\, q} \; \bar{a}^{\dagger\, 2k-2j-q} \\ &= \tfrac{1}{\sqrt{2}^{2k+1}} \sum_{p=0}^{2j+1} \sum_{q=0}^{2k-2j} \binom{2j+1}{p} \binom{2k-2j}{q} \\ &\quad (-1)^q \; a^{\dagger\, p+q} \; \bar{a}^{\dagger\, 2k+1-p-q} . \end{aligned}$$

The multi-photon states expressed in the complementary basis are as follows

$$\begin{aligned} |\Phi_k^P\rangle = &\tfrac{1}{\sqrt{\mathcal{N}_k^P}} \tfrac{1}{\sqrt{2}^{2k+1}} \sum_{\substack{j=0 \\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{\infty} \sum_{p=0}^{2j+1} \sum_{q=0}^{2k-2j} \qquad (33) \\ &\binom{k}{j} \binom{2j+1}{p} \binom{2k-2j}{q} \\ &(-1)^q \sqrt{(p+q)! \, (2k+1-p-q)!} \\ &|p+q, 2k+1-p-q\rangle_{\frac{\pi}{2}} , \end{aligned}$$

$$\begin{aligned} |\bar{\Phi}_k^P\rangle = &\tfrac{1}{\sqrt{\bar{\mathcal{N}}_k^P}} \tfrac{1}{\sqrt{2}^{2k+1}} \sum_{\substack{j=0 \\ \mathrm{C}(\sigma=2k+1,\Delta=4j-1-2k)}}^{\infty} \sum_{p=0}^{2j} \sum_{q=0}^{2k+1-2j} \qquad (34) \\ &\binom{k}{j} \binom{2j}{p} \binom{2k+1-2j}{q} \\ &(-1)^{1-q} \sqrt{(p+q)! \, (2k+1-p-q)!} \\ &|p+q, 2k+1-p-q\rangle_{\frac{\pi}{2}} . \end{aligned}$$

Due to the orthogonality of Fock states, A^P takes the form of a convex sum

$$\begin{gathered} \langle \Phi_k^P | \mathcal{O}^{(M)}(\tfrac{\pi}{2}) | \bar{\Phi}_{k'}^P \rangle \sim \\ \langle p+q, 2k+1-p-q | \tilde{n}, \tilde{m} \rangle \langle \tilde{n}, \tilde{m} | p'+q', 2k'+1-p'-q' \rangle \\ \downarrow \\ p+q = p'+q', k = k' \end{gathered}$$

and

$$\begin{aligned} A_k^P = &\tfrac{1}{\sqrt{\mathcal{N}_k^P \, \bar{\mathcal{N}}_k^P}} \tfrac{1}{2^{2k+1}} \sum_{\substack{j=0 \\ \mathrm{C}(\sigma=2k+1,\Delta=4j+1-2k)}}^{k} \sum_{p=0}^{2j+1} \sum_{q=0}^{2k-2j} \qquad (35) \\ &\sum_{\substack{j'=0 \\ \mathrm{C}(\sigma=2k+1,\Delta=4j'-1-2k)}}^{k} \sum_{p'=0}^{2j'} \sum_{q'=0}^{2k+1-2j'} \\ &\binom{k}{j} \binom{2j+1}{p} \binom{2k-2j}{q} (-1)^q \\ &(p+q)! \, (2k+1-p-q)! \, \mathrm{sgn}(2(p+q)-2k-1) \\ &\binom{k}{j'} \binom{2j'}{p'} \binom{2k+1-2j'}{q'} (-1)^{1-q'} \, \delta_{p+q=p'+q'} . \end{aligned}$$

The Bell parameter takes the form

$$\begin{aligned} B &= 2 \cos\alpha \, V(0) + 2 \sin\alpha \, A(\pi/2), \\ B &= 2 \cos\alpha \sum_{k=0}^{\infty} \left(\beta_k^P\right)^2 V_k^P + 2 \sin\alpha \sum_{k=0}^{\infty} \left(\beta_k^P\right)^2 A_k^P , \\ B &= \sum_{k=0}^{\infty} \left(\beta_k^P\right)^2 B_k^P , \qquad (36) \\ B_k^P &= 2 \sqrt{\left(V_k^P\right)^2 + \left(A_k^P\right)^2} . \qquad (37) \end{aligned}$$

[1] M. Arndt, T. Juffmann, an V. Vedral, HFSP Journal **3**, 386 (2009); M. Arndt *et al*, Nature **401**, 680 (1999); B. Brezger *et al*, Phys. Rev. Lett. **88**, 100404 (2002).
[2] J. S. Bell, Physics **1**, 195 (1965).
[3] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
[4] M. A. Rowe *et al* Nature **409**, 791-794 (2001); S. Gröblacher *et al*, Nature **446**, 871 (2007); N. Brunner, N. Gisin, V. Scarani, and C. Simon, Phys. Rev. Lett. **98**, 220403 (2007).

[5] F. De Martini, F. Sciarrino, and C. Vitelli, Phys. Rev. Lett. **100**, 253601 (2008).

[6] T. Sh. Iskhakov, M. V. Chekhova, G. O. Rytikov, and G. Leuchs, Phys. Rev. Lett. **106**, 113602 (2011).

[7] C. Vitelli, N. Spagnolo, L. Toffoli, F. Sciarrino, and F. De Martini, Phys. Rev. A **81**, 032123 (2010).

[8] M. Stobińska, P. Sekatski, A. Buraczewski, N. Gisin, and G. Leuchs, Phys. Rev. A **84**, 034104 (2011).

[9] P. Sekatski, N. Brunner, C. Branciard, N. Gisin and C. Simon, Phys. Rev. Lett. **103**, 113601 (2009).

[10] M. Stobińska, P. Horodecki, A. Buraczewski, R. W. Chhajlany, R. Horodecki, G. Leuchs, arXiv:0909.1545.

[11] C. Vitelli, N. Spagnolo, F. Sciarrino, and F. De Martini, Phys. Rev. A **82**, 062319 (2010).

[12] M. Stobińska, F. Töppel, P. Sekatski, A. Buraczewski, M. Żukowski M. V. Chekhova, N. Gisin, and G. Leuchs, Phys. Rev. A **86**, 063823 (2012).

[13] P. Sekatski, B. Sanguinetti, E. Pomarico, N. Gisin, and C. Simon, Phys. Rev. A **82**, 053814 (2010).

[14] F. De Martini, F. Sciarrino, and Ch. Vitelli, Phys. Rev. Lett. **100**, 253601 (2008).

[15] T. Iskhakov, M. V. Chekhova, and G. Leuchs, Phys. Rev. Lett. **102**, 183602 (2009).